Probability Theory & Mathematical Statistics

Moment and Deviation

尹一通 Nanjing University, 2026 Spring

Moments and Deviations



Markov's Inequality

(马尔可夫不等式)



- <u>**Markov's inequality**</u>: Let X be a *nonnegative-valued* random variable. Then,

$$\text{for any } a > 0, \qquad \Pr(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$$

- **Proof** (by indicator): Let $I = I(X \geq a)$. Since $X \geq 0$ and $a > 0$, we have

$$I = I(X \geq a) \leq \left\lfloor \frac{X}{a} \right\rfloor \leq \frac{X}{a}.$$

Therefore, $\Pr(X \geq a) = \mathbb{E}[I] \leq \mathbb{E}\left[\frac{X}{a}\right] = \frac{\mathbb{E}[X]}{a}$



Markov's Inequality

(马尔可夫不等式)



- <u>**Markov's inequality**</u>: Let X be a *nonnegative-valued* random variable. Then,

$$\text{for any } a > 0, \qquad \Pr(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$$

- **Proof** (by total expectation):

$$\mathbb{E}[X] = \underbrace{\mathbb{E}[X \mid X \geq a]}_{(X \geq a \text{ is possible})} \cdot \Pr(X \geq a) + \underbrace{\mathbb{E}[X \mid X < a]}_{(X \text{ is nonnegative})} \cdot \Pr(X < a)$$

$$\geq a \cdot \Pr(X \geq a) + 0 \cdot \Pr(X < a) \qquad = a \cdot \Pr(X \geq a)$$

$$\implies \Pr(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$$

Markov's Inequality

(马尔可夫不等式)



- <u>**Markov's inequality**</u>: Let X be a *nonnegative-valued* random variable. Then,

$$\text{for any } a > 0, \qquad \Pr(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$$

- **Corollary**: for any $c > 1$, $\quad \Pr(X \geq c\mathbb{E}[X]) \leq 1/c$

- **Tight in the worst case**: $\forall c > 1, \forall \mu \in \mathbb{R}$, $\exists$ nonnegative X with $\mathbb{E}[X] = \mu$, such that $\Pr(X \geq c\mu) = 1/c$

- **Lower tail variant** (sometimes called <u>*reverse Markov's inequality*</u>):

  $\Pr(X \leq a) \leq (u - \mathbb{E}[X])/(u - a)$ requires X to have bounded range $X \leq u$

From Las Vegas to Monte Carlo

-

<u>Monte Carlo algorithm</u>: randomized algorithms that are correct by chance
-

<u>Las Vegas algorithm</u>: randomized algorithms that always give correct result upon termination (but may run for a random period of time before termination)

- If there is a Las Vegas algorithm $\mathscr{A}$ with expected running time at most $t(n)$ for any input of size n ($\mathscr{A}$ has worst-case expected time complexity $t(n)$):

Algorithm $\mathscr{B}$:

simulate algorithm $\mathscr{A}$ up to $\lceil t(n)/\epsilon \rceil$ steps;
if algorithm $\mathscr{A}$ terminates
    return the output of $\mathscr{A}$;
else return an arbitrary answer;

- Algorithm $\mathscr{B}$ is a Monte Carlo algorithm s.t.
  - $\mathscr{B}$ has worst-case running time $\leq \lceil t(n)/\epsilon \rceil$
  - $\mathscr{B}$ is correct with probability at least $1 - \epsilon$
    (by Markov inequality)

Cliques in Random Graph

- $G(n,p)$: between every pair u, v among n vertices, an edge is added i.i.d. with prob. p
- Fix a constant integer $k \geq 3$. Let X be the number of k-**cliques** (K_k) in $G \sim G(n,p)$.
- For every distinct $S \subseteq \in [n]$ of size $|S| = k$, let $I_S = I(K_S \subseteq G)$. Then:
  - $\mathbb{E}[I_S] = \Pr(K_S \subseteq G) = p^{\binom{k}{2}}$
  - $X = \sum_{S \in \binom{[n]}{k}} I_S$



- **Linearity of expectation:** $\mathbb{E}[X] = \binom{n}{k} p^{\binom{k}{2}} \leq n^k p^{k(k-1)/2} = o(1)$ for $p = o\left(n^{-2/(k-1)}\right)$
- **Markov's inequality:** $\Pr(X \geq 1) \leq \mathbb{E}[X] = o(1) \Longrightarrow \Pr(X = 0) = 1 - o(1)$
  $\Longrightarrow$ If $p = o\left(n^{-2/(k-1)}\right)$, then $G(n,p)$ is K_k-free **a.a.s.** (asymptotically almost surely)

Generalized Markov's Inequality

- Let X be a random variable and $f : \mathbb{R} \to \mathbb{R}_{\geq 0}$ a nonnegative-valued function.

$$\text{For any } a > 0, \qquad \Pr(f(X) \geq a) \leq \frac{\mathbb{E}[f(X)]}{a}$$

- **Proof**: Apply the Markov's inequality to the random variable $Y = f(X)$.
- **Applications**: useful if $f(X)$ can "*extract*" useful information about X
  - **Chebyshev's inequality**, k**th moment method**: $f(X)$ extracts the kth moment
  - **Chernoff-Hoeffding bounds, Bernstein inequalities**: $f(X)$ extracts all moments

Deviation Inequality



- Let X be a random variable with **mean** $\mu = \mathbb{E}[X]$. For $a > 0$

$$\Pr(|X - \mu| \geq a) \leq ?$$

- Applying **Markov's inequality** to $Y = |X - \mu|$ gives us

$$\Pr(|X - \mu| \geq a) \leq \frac{\mathbb{E}[|X - \mu|]}{a}$$

difficult to calculate

- Alternatively, we may apply **Markov's inequality** to $Y = (X - \mu)^2$

$$\Pr(|X - \mu| \geq a) = \Pr((X - \mu)^2 \geq a^2) \leq \frac{\mathbb{E}[(X - \mu)^2]}{a^2}$$

Variance

(2nd central moment)

Variance (方差) and Moments (矩)

- For integer $k > 0$, the <u>kth moment</u> (k阶矩) of a random variable X is $\mathbb{E}[X^k]$, and the <u>kth central moment</u> (k阶中心矩) of X is $\mathbb{E}[(X - \mathbb{E}[X])^k]$.

- Sometimes, a random variable X is called **centralized** (中心化的) if $\mathbb{E}[X] = 0$. A random variable X can be centralized by $Y = X - \mathbb{E}[X]$.

- The <u>variance</u> (方差) of a random variable X is its 2nd central moment:

$$\mathbf{Var}[X] = \mathbb{E}\left[(X - \mathbb{E}[X])^2\right]$$

  and the <u>**standard deviation**</u> (标准差) of X is $\sigma = \sigma[X] = \sqrt{\mathbf{Var}[X]}$

Chebyshev's Inequality

(切比雪夫不等式)



- <u>Chebyshev's inequality</u>: Let X be a random variable. For any $a > 0$,

$$\Pr(|X - \mathbb{E}[X]| \geq a) \leq \frac{\mathbf{Var}[X]}{a^2}$$

- **Proof**: Apply Markov's inequality to $Y = (X - \mathbb{E}[X])^2$.

- **Corollary**: For standard deviation $\sigma = \sqrt{\mathbf{Var}[X]}$, for any $k \geq 1$,

$$\Pr(|X - \mathbb{E}[X]| \geq k\sigma) \leq \frac{1}{k^2}$$

Median and Mean

- The <u>median</u> (中位数) of random variable X is defined to be any value m s.t.:

$$\Pr(X \le m) \ge 1/2 \quad \text{and} \quad \Pr(X \ge m) \ge 1/2$$

- The expectation $\mu = \mathbb{E}[X]$ is the value that minimizes

$$\mathbb{E}[(X-\mu)^2]$$

- **Proof**: $f(x) = \mathbb{E}[(X-x)^2] = \mathbb{E}[X^2] - 2x\mathbb{E}[X] + x^2$ is convex and has $f'(\mu) = 0$

- The median m is the value that minimizes

$$\mathbb{E}[\,|X-m|\,]$$

- **Proof**: By symmetry, suppose non-median $y > m$ so that $\Pr(X \ge y) < 1/2$.

$$\mathbb{E}[\,|X-y| - |X-m|\,] = (y-m)\Pr(X \le m) + \sum_{m<x<y} (m+y-2x)\Pr(X=x) + (m-y)\Pr(X \ge y)$$

$$> (y-m)\Pr(X \le m) + \sum_{m<x<y} (m-y)\Pr(X=x) + (m-y)\Pr(X \ge y) \ge (y-m)/2 + (m-y)/2 = 0$$

Median and Mean



- If X is a random variable with finite expectation μ, median m, and standard deviation σ, then

$$|\mu - m| \leq \sigma$$

- **Proof**: $|\mu - m| = |\mathbb{E}[X] - m| = |\mathbb{E}[X - m]|$

$$\leq \mathbb{E}[\,|X - m|\,] \quad \text{(Jensen's inequality)}$$

$$\leq \mathbb{E}[\,|X - \mu|\,] \quad \text{(the median } m \text{ minimizes } \mathbb{E}[\,|X - m|\,]\text{)}$$

$$= \mathbb{E}\left[\sqrt{(X-\mu)^2}\right] \leq \sqrt{\mathbb{E}\left[(X-\mu)^2\right]} = \sigma \quad \text{(Jensen's inequality)}$$

Variance

Calculation of Variance

$$\mathbf{Var}[X] = \mathbb{E}\left[(X - \mathbb{E}[X])^2\right] = \mathbb{E}[X^2] - \mathbb{E}[X]^2$$

- **Proof**: $\mathbf{Var}[X] = \mathbb{E}\left[(X - \mathbb{E}[X])^2\right]$

$$= \mathbb{E}\left[X^2 - 2\mathbb{E}[X]X + \mathbb{E}[X]^2\right]$$

$$= \mathbb{E}[X^2] - 2\mathbb{E}[X]\mathbb{E}[X] + \mathbb{E}[X]^2$$

$$= \mathbb{E}[X^2] - \mathbb{E}[X]^2$$

- X is constant ***a.s.*** $(\Pr(X = \mathbb{E}[X]) = 1) \iff \mathbb{E}[X^2] = \mathbb{E}[X]^2 \iff \mathbf{Var}[X] = 0$

Variance of Linear Function

- For random variables X, Y and real number $a \in \mathbb{R}$:
  - $\mathbf{Var}[a] = 0$
  - $\mathbf{Var}[X + a] = \mathbf{Var}[X]$ (variance is a central moment)
  - $\mathbf{Var}[aX] = a^2\mathbf{Var}[X]$ (variance is quadratic)
  - $\mathbf{Var}[X + Y] = \mathbf{Var}[X] + \mathbf{Var}[Y] + 2(\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y])$
- **Proof**: All can be verified through $\mathbf{Var}[X] = \mathbb{E}[X^2] - \mathbb{E}[X]^2$.

Covariance (协方差)

- The covariance (协方差) of two random variables X and Y is

$$\mathbf{Cov}(X, Y) = \mathbb{E}\left[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])\right] = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$$

- **Properties**: $\mathbf{Var}[X] = \mathbf{Cov}(X, X)$
  - *Symmetric*: $\mathbf{Cov}(X, Y) = \mathbf{Cov}(Y, X)$
  - *Distributive*: $\mathbf{Cov}(X + Y, Z) = \mathbf{Cov}(X, Z) + \mathbf{Cov}(Y, Z)$
    $\mathbf{Cov}(aX, Y) = a\mathbf{Cov}(X, Y)$

- If X and Y are **independent** then

$$\mathbf{Cov}(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y] = 0$$

Covariance of Independent Variables

- If random variables X and Y are **independent**, then

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

- If random variables $X_1, X_2, \ldots, X_n$ are **mutually independent**, then

$$\mathbb{E}\left[\prod_{i=1}^{n} X_i\right] = \mathbb{E}\left[\prod_{i=1}^{n-1} X_i\right] \cdot \mathbb{E}[X_n] = \prod_{i=1}^{n} \mathbb{E}[X_i]$$

Proof: By change of variable (*LOTUS*)

$$\mathbb{E}[XY] = \sum_{x,y} xy \Pr(X = x \cap Y = y) = \sum_{x,y} xy \Pr(X = x) \Pr(Y = y)$$

$$= \left(\sum_{x} x \Pr(X = x)\right)\left(\sum_{y} y \Pr(Y = y)\right) = \mathbb{E}[X]\mathbb{E}[Y]$$

Expectation of Product

- For random variables X and Y:

$$\text{if } X \text{ and } Y \text{ independent, then } \mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

- **(Cauchy-Schwarz)**

$$\mathbb{E}[XY]^2 \leq \mathbb{E}[X^2]\mathbb{E}[Y^2]$$

Proof:

$$0 \leq \mathbb{E}\left[\left(X - \frac{\mathbb{E}[XY]}{\mathbb{E}[Y^2]}Y\right)^2\right] = \mathbb{E}[X^2] - \frac{\mathbb{E}[XY]^2}{\mathbb{E}[Y^2]}$$

$$\implies \mathbb{E}[X^2]\mathbb{E}[Y^2] \geq \mathbb{E}[XY]^2$$

- **(Hölder)** for any $p, q > 0$ satisfying $\frac{1}{p} + \frac{1}{q} = 1$

$$\mathbb{E}[XY] \leq \mathbb{E}[|X|^p]^{1/p}\mathbb{E}[|Y|^q]^{1/q}$$

Correlation (相关性)



- The <u>covariance</u> (协方差) of two random variables X and Y is

$$\mathbf{Cov}(X, Y) = \mathbb{E}\left[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])\right] = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$$

- The <u>correlation coefficient</u> (相关性系数) of X and Y is

$$\rho(X, Y) = \frac{\mathbf{Cov}(X, Y)}{\sqrt{\mathbf{Var}[X] \cdot \mathbf{Var}[Y]}}$$

$\in [-1,1]$ by Cauchy-Schwarz

Apply Cauchy-Schwarz to:

$$U = \frac{X - \mathbb{E}[X]}{\sigma(X)}, V = \frac{Y - \mathbb{E}[Y]}{\sigma(Y)}$$

- Two random variables X and Y are called <u>uncorrelated</u> if $\mathbf{Cov}(X, Y) = 0$

- X and Y are **uncorrelated** means:
  - $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$
  - $\mathbf{Var}[X + Y] = \mathbf{Var}[X] + \mathbf{Var}[Y]$

Uncorrelated but not independent:

- $X \sim U[-1,1]$, $Y = X^2$;
- $X = Z_1 + Z_2$, $Y = Z_1 - Z_2$,
  where $Z_1, Z_2 \sim$ Bernoulli$(1/2)$ i.i.d.

Variance of Sum

- For random variables X, Y:

$$\mathbf{Var}[X+Y] = \mathbf{Var}[X] + \mathbf{Var}[Y] + 2\mathbf{Cov}(X, Y)$$

- For random variables $X_1, X_2, \ldots, X_n$:

$$\mathbf{Var}\left[\sum_{i=1}^{n} X_i\right] = \sum_{i=1}^{n} \mathbf{Var}[X_i] + \sum_{i \neq j} \mathbf{Cov}(X_i, X_j)$$

- For pairwise independent $X_1, X_2, \ldots, X_n$:

$$\mathbf{Var}\left[\sum_{i=1}^{n} X_i\right] = \sum_{i=1}^{n} \mathbf{Var}[X_i]$$

Variance of Indicator



- For **Bernoulli random variable** $X \in \{0,1\}$ with parameter p

$$X^2 = X \implies \mathbb{E}[X^2] = \mathbb{E}[X] = p$$

$$\mathbf{Var}[X] = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = p - p^2 = p(1-p)$$

- For the **indicator** random variable $X = I(A)$ of event A:

$$\mathbf{Var}[X] = \Pr(A)(1 - \Pr(A)) = \Pr(A)\Pr(A^c)$$

Variance of Discrete Uniform Distribution

- For integers $a \leq b$, let X be chosen from $[a, b] = \{a, a+1, \ldots, b\}$ **u.a.r.**
  - $\mathbb{E}[X] = \sum_{k=a}^{b} \frac{k}{b-a+1} = \frac{a+b}{2}$
  - $\mathbb{E}[X^2] = \sum_{k=a}^{b} \frac{k^2}{b-a+1} = \frac{2b^2 + 2ab + 2a^2 + b - a}{6}$
  - $\mathbf{Var}[X] = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \frac{(b-a)(b-a+2)}{12}$

Poisson Distribution

- For **Poisson random variable** $X \sim \text{Pois}(\lambda)$, recall $\mathbb{E}[X] = \lambda$, and

$$\begin{aligned}
\mathbb{E}[X^2] &= \sum_{k \geq 0} k^2 \frac{\mathrm{e}^{-\lambda}\lambda^k}{k!} = \sum_{k \geq 1} k \frac{\mathrm{e}^{-\lambda}\lambda^k}{(k-1)!} \\
&= \sum_{k \geq 0} (k+1) \frac{\mathrm{e}^{-\lambda}\lambda^{k+1}}{k!} = \lambda \sum_{k \geq 0} (k+1) \frac{\mathrm{e}^{-\lambda}\lambda^k}{k!} \\
&= \lambda \mathbb{E}[X+1] = \lambda(\mathbb{E}[X]+1) = \lambda(\lambda+1)
\end{aligned}$$

$$\mathbf{Var}[X] = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \lambda(\lambda+1) - \lambda^2 = \lambda$$

Geometric Distribution (几何分布)

- For **geometric random variable** $X \sim \text{Geo}(p)$, recall $\mathbb{E}[X] = 1/p$, and

$$\mathbb{E}[X^2] = \sum_{k \geq 1} k^2 (1-p)^{k-1} p = (2-p)p^{-2}$$

$$\mathbf{Var}[X] = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = (2-p)p^{-2} - p^{-2} = (1-p)/p^2$$

- **Total expectation:** $\mathbb{E}[X^2] = \mathbb{E}[X^2 \mid X > 1] \cdot (1-p) + \mathbb{E}[X^2 \mid X = 1] \cdot p$

$$= \mathbb{E}[((X-1)+1)^2 \mid X > 1] \cdot (1-p) + p$$

(memoryless) $$= \mathbb{E}[(X+1)^2] \cdot (1-p) + p$$

$$= (1-p)\mathbb{E}[X^2] + 2(1-p)/p + 1$$

$$\implies \mathbb{E}[X^2] = (2-p)/p^2 \implies \mathbf{Var}[X] = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = (1-p)/p^2$$

Binomial Distribution (二项分布)

- For **binomial random variable** $X \sim \text{Bin}(n, p)$, recall $\mathbb{E}[X] = np$, and

$$\mathbf{Var}[X] = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \sum_{k=0}^{n} k^2 \binom{n}{k} p^k (1-p)^{n-k} - (np)^2$$

- **Observation**: $X \sim \text{Bin}(n, p)$ can be expressed as $X = X_1 + \cdots + X_n$, where $X_1, \ldots, X_n$ are i.i.d. Bernoulli random variables with parameter p

- For **mutually independent** $X_1, \ldots, X_n$:

$$\mathbf{Var}\,[X] = \sum_{i=1}^{n} \mathbf{Var}[X_i] \; = np(1-p)$$

Negative Binomial Distribution (负二项分布)

- For **negative binomial random variable** X with parameters r, p

$$\mathbf{Var}[X] = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \sum_{k \geq 1} k^2 \binom{k+r-1}{k} (1-p)^k p^r - r^2(1-p)^2/p^2$$

- **Observation**: X can be expressed as $X = (X_1 - 1) + \cdots + (X_r - 1)$, where $X_1, \ldots, X_r$ are i.i.d. geometric random variables with parameter p

- For **mutually independent** $X_1, \ldots, X_r$:

$$\mathbf{Var}\,[X] = \sum_{i=1}^{r} \mathbf{Var}[X_i - 1] = \sum_{i=1}^{r} \mathbf{Var}[X_i] = \frac{r(1-p)}{p^2}$$

Chebyshev (Чебышёв)'s Inequality



Chebyshev's Inequality

(切比雪夫不等式)



- <u>Chebyshev's inequality</u>: Let X be a random variable. For any $a > 0$,

$$\Pr(|X - \mathbb{E}[X]| \geq a) \leq \frac{\mathbf{Var}[X]}{a^2}$$

- **Corollary**: For standard deviation $\sigma = \sqrt{\mathbf{Var}[X]}$, for any $k \geq 1$,

$$\Pr(|X - \mathbb{E}[X]| \geq k\sigma) \leq \frac{1}{k^2}$$

- **Tight in the worst case**: $\forall k \geq 1$, $\forall \mu \in \mathbb{R}$ and $\forall \sigma > 0$, $\exists X$ with $\mathbb{E}[X] = \mu$ and $\mathbf{Var}[X] = \sigma^2$ such that $\Pr(|X - \mu| \geq k\sigma) = 1/k^2$

Mean Trick

- Let $X_1, \ldots, X_n$ be *i.i.d.* random variables with $\mathbb{E}[X_i] = \mu$ and $\mathbf{Var}[X_i] = \sigma^2$.
- Empirical mean: $\overline{X} = \frac{1}{n}\sum_{i=1}^{n} X_i$

$$\mathbb{E}[\overline{X}] = \frac{1}{n}\sum_{i=1}^{n}\mathbb{E}[X_i] = \mu \text{ and } \mathbf{Var}[\overline{X}] = \frac{1}{n^2}\sum_{i=1}^{n}\mathbf{Var}[X_i] = \frac{\sigma^2}{n}$$

- **Chebyshev's inequality**:

$$\Pr(|\overline{X} - \mu| \geq \epsilon\mu) \leq \frac{\mathbf{Var}[\overline{X}]}{\epsilon^2\mu^2} = \frac{\sigma^2}{\epsilon^2\mu^2 n} \leq \delta \text{ if } n \geq \frac{\sigma^2}{\epsilon^2\mu^2\delta}$$

Balls into Bins



- Throwing n balls into n bins, the most heavily loaded bin receives $\Theta\left(\frac{\ln n}{\ln \ln n}\right)$ balls ***a.a.s.*** (***asymptotically almost surely***, with probability $1 - o(1)$)

- **Proof of $O\left(\frac{\ln n}{\ln \ln n}\right)$ upper bound:** Choose $k = 3 \ln n / \ln \ln n$

$$\Pr(\text{max-load} \geq k) = \Pr(\exists \text{ bin recives } \geq k \text{ balls}) \leq n \Pr(\text{bin-1 recives } \geq k \text{ balls}) \leq \frac{1}{n}$$

(union bound)

$$\Pr(\text{bin-1 recives } \geq k \text{ balls})$$

$$= \Pr\left(\exists S \in \binom{\{n \text{ balls}\}}{k} \text{ s.t. all balls in } S \text{ are assigned to bin-1} \right)$$

$$\leq \binom{n}{k} \frac{1}{n^k} \leq \left(\frac{\mathrm{e}n}{k}\right)^k \frac{1}{n^k} \leq \left(\frac{\mathrm{e}}{k}\right)^k \leq \frac{1}{n^2}$$

(union bound)

Balls into Bins



- Throwing n balls into n bins, the most heavily loaded bin receives $\Theta\left(\frac{\ln n}{\ln \ln n}\right)$ balls ***a.a.s.*** (***asymptotically almost surely***, with probability $1 - o(1)$)

- **Proof of $\Omega\left(\frac{\ln n}{\ln \ln n}\right)$ lower bound:** Choose $k = \ln n / 2 \ln \ln n$

Let $I_i = I(\text{bin-}i \text{ recives extactly } k \text{ balls})$ and let $X = \sum_{i=1}^{n} I_i$

$$\Pr(\text{max-load} < k) = \Pr(\text{every bin recives } < k \text{ balls}) \leq \Pr(X = 0)$$

$$\leq \Pr(|X - \mathbb{E}[X]| = \mathbb{E}[X]) \leq \frac{\mathbf{Var}[X]}{\mathbb{E}[X]^2}$$ (Chebyshev)

(use $\binom{n}{k} \geq \left(\frac{n}{k}\right)^k$)

Define $p = \mathbb{E}[I_i] = \binom{n}{k} n^{-k} \left(1 - \frac{1}{n}\right)^{n-k}$, then $\mathbb{E}[X] = np = \binom{n}{k} n^{1-k} \left(1 - \frac{1}{n}\right)^{n-k} \geq \frac{n}{\mathrm{e}k^k} = \omega(1)$

(binomial distribution)

Balls into Bins



- Throwing n balls into n bins, the most heavily loaded bin receives $\Theta\left(\frac{\ln n}{\ln \ln n}\right)$ balls ***a.a.s.*** (***asymptotically almost surely***, with probability $1 - o(1)$)

- **Proof of $\Omega\left(\frac{\ln n}{\ln \ln n}\right)$ lower bound:** Choose $k = \ln n / 2 \ln \ln n$

Let $I_i = I(\text{bin-}i \text{ recives extactly } k \text{ balls})$ and let $X = \sum_{i=1}^{n} I_i$

Define:

$$p = \mathbb{E}[I_i] = \binom{n}{k} n^{-k} \left(1 - \frac{1}{n}\right)^{n-k}$$

$$\Pr(\text{max-load} < k) \leq \frac{\mathbf{Var}[X]}{\mathbb{E}[X]^2} \leq \frac{1}{np} + \frac{\sum_{i \neq j} \mathbb{E}[I_i I_j] - (np)^2}{(np)^2} = o(1) + \frac{\sum_{i \neq j} \mathbb{E}[I_i I_j] - (np)^2}{(np)^2}$$

$\mathbb{E}[X] = np = \omega(1)$ and $\mathbf{Var}[X] = \sum_{i=1}^{n} \mathbf{Var}[I_i] + \sum_{i \neq j} \mathbf{Cov}(I_i, I_j) = \sum_{i,j} (\mathbb{E}[I_i I_j] - \mathbb{E}[I_i]\mathbb{E}[I_j]) = \sum_{i,j} (\mathbb{E}[I_i I_j] - p^2)$

$$= \sum_{i=1}^{n} \mathbb{E}[I_i^2] + \sum_{i \neq j} \mathbb{E}[I_i I_j] - (np)^2 = \sum_{i=1}^{n} \mathbb{E}[I_i] + \sum_{i \neq j} \mathbb{E}[I_i I_j] - (np)^2 = np + \sum_{i \neq j} \mathbb{E}[I_i I_j] - (np)^2$$

Balls into Bins



- Throwing n balls into n bins, the most heavily loaded bin receives $\Theta\left(\frac{\ln n}{\ln \ln n}\right)$ balls ***a.a.s.*** (***asymptotically almost surely***, with probability $1 - o(1)$)

- **Proof of $\Omega\left(\frac{\ln n}{\ln \ln n}\right)$ lower bound:** Choose $k = \ln n / 2 \ln \ln n$

Let $I_i = I(\text{bin-}i \text{ recives extactly } k \text{ balls})$ and let $X = \sum_{i=1}^{n} I_i$

Define:

$$p = \mathbb{E}[I_i] = \binom{n}{k} n^{-k} \left(1 - \frac{1}{n}\right)^{n-k}$$

$$\Pr(\text{max-load} < k) \leq \frac{\mathbf{Var}[X]}{\mathbb{E}[X]^2} \leq o(1) + \frac{\sum_{i \neq j} \mathbb{E}[I_i I_j] - (np)^2}{(np)^2} = o(1)$$

$$\mathbb{E}[I_i I_j] = \Pr(\text{both bins-}i, j \text{ recive extactly } k \text{ balls}) = \binom{n}{k, k, n-2k} n^{-2k} \left(1 - \frac{2}{n}\right)^{n-2k} \leq (1 + o(1)) p^2$$

(multinomial distribution)

Cliques in Random Graph (revisited)

- Fix a constant integer $k \geq 3$. Let X be the number of k-**cliques** (K_k) in $G \sim G(n, p)$.

- For every distinct $S \subseteq \in [n]$ of size $|S| = k$, let $I_S = I(K_S \subseteq G)$. Then:

$$X = \sum_{S \in \binom{[n]}{k}} I_S \text{ and } \mathbb{E}[I_S] = \Pr(K_S \subseteq G) = p^{\binom{k}{2}}$$



- **Linearity of expectation:** $\mathbb{E}[X] = \binom{n}{k} p^{\binom{k}{2}} = \Theta\left(n^k p^{\binom{k}{2}}\right)$

$$\mathbb{E}[X] = \Theta\left(n^k p^{\binom{k}{2}}\right) = \begin{cases} o(1) & \text{if } p = o\left(n^{-2/(k-1)}\right) \\ \omega(1) & \text{if } p = \omega\left(n^{-2/(k-1)}\right) \end{cases}$$

$$\xRightarrow{\text{(Markov)}} \Pr(X \geq 1) = o(1)$$

$$\xRightarrow{?} \Pr(X \geq 1) = 1 - o(1)$$

Cliques in Random Graph (revisited)

- Fix a constant integer $k \geq 3$. Let X be the number of k-**cliques** (K_k) in $G \sim G(n,p)$.

- For every distinct $S \subseteq \in [n]$ of size $|S| = k$, let $I_S = I(K_S \subseteq G)$. Then:

$$X = \sum_{S \in \binom{[n]}{k}} I_S \text{ and } \mathbb{E}[X] = \Theta\left(n^k p^{\binom{k}{2}}\right) = \begin{cases} o(1) & \text{if } p = o\left(n^{-2/(k-1)}\right) \\ \omega(1) & \text{if } p = \omega\left(n^{-2/(k-1)}\right) \end{cases}$$

- **Chebyshev**: $\Pr(X = 0) \leq \Pr(|X - \mathbb{E}[X]| \geq \mathbb{E}[X]) \leq \dfrac{\mathbf{Var}[X]}{\mathbb{E}[X]^2} \leq \dfrac{1}{\mathbb{E}[X]} + \dfrac{\sum_{S \neq T} \mathbb{E}[I_S I_T]}{\mathbb{E}[X]^2}$

$$\mathbf{Var}[X] = \sum_{S \in \binom{[n]}{k}} \mathbf{Var}[I_S] + \sum_{\substack{S \neq T \\ S,T \in \binom{[n]}{k}}} \mathbf{Cov}(I_S, I_T) = \sum_{S \in \binom{[n]}{k}} (\mathbb{E}[I_S^2] - \mathbb{E}[I_S]^2) + \sum_{\substack{S \neq T \\ S,T \in \binom{[n]}{k}}} (\mathbb{E}[I_S I_T] - \mathbb{E}[I_S]\mathbb{E}[I_T])$$

$$= \sum_S (\mathbb{E}[I_S] - \mathbb{E}[I_S]^2) + \sum_{S \neq T} (\mathbb{E}[I_S I_T] - \mathbb{E}[I_S]\mathbb{E}[I_T])$$

$$\leq \mathbb{E}[X] + \sum_{S \neq T} \mathbb{E}[I_S I_T]$$

Cliques in Random Graph (revisited)

- Fix a constant integer $k \geq 3$. Let X be the number of k-**cliques** (K_k) in $G \sim G(n, p)$.

- For every distinct $S \subseteq \in [n]$ of size $|S| = k$, let $I_S = I(K_S \subseteq G)$. Then:

$$X = \sum_{S \in \binom{[n]}{k}} I_S \text{ and } \mathbb{E}[X] = \Theta\left(n^k p^{\binom{k}{2}}\right) = \begin{cases} o(1) & \text{if } p = o\left(n^{-2/(k-1)}\right) \\ \omega(1) & \text{if } p = \omega\left(n^{-2/(k-1)}\right) \end{cases}$$

- **Chebyshev**: $\Pr(X = 0) \leq \Pr(|X - \mathbb{E}[X]| \geq \mathbb{E}[X]) \leq \dfrac{\mathbf{Var}[X]}{\mathbb{E}[X]^2} \leq \dfrac{1}{\mathbb{E}[X]} + \dfrac{\sum_{S \neq T} \mathbb{E}[I_S I_T]}{\mathbb{E}[X]^2}$

$$\mathbb{E}[I_S I_T] = \Pr((K_S \cup K_T) \subseteq G) = p^{2\binom{k}{2} - \binom{|S \cap T|}{2}}$$

$$\sum_{\substack{S \neq T \\ S, T \in \binom{[n]}{k}}} \mathbb{E}[I_S I_T] = \sum_{\ell=2}^{k-1} \sum_{\substack{|S \cap T| = \ell \\ S, T \in \binom{[n]}{k}}} \mathbb{E}[I_S I_T] = \sum_{\ell=2}^{k-1} \binom{n}{2k - \ell} \cdot O(1) \cdot p^{2\binom{k}{2} - \binom{\ell}{2}} = O\left(n^{2k} p^{2\binom{k}{2}} \sum_{\ell=2}^{k-1} n^{-\ell} p^{-\binom{\ell}{2}}\right)$$

Cliques in Random Graph (revisited)

- Fix a constant integer $k \geq 3$. Let X be the number of k-**cliques** (K_k) in $G \sim G(n,p)$.

- For every distinct $S \subseteq \in [n]$ of size $|S| = k$, let $I_S = I(K_S \subseteq G)$. Then:

$$X = \sum_{S \in \binom{[n]}{k}} I_S \text{ and } \mathbb{E}[X] = \Theta\left(n^k p^{\binom{k}{2}}\right) = \begin{cases} o(1) & \text{if } p = o\left(n^{-2/(k-1)}\right) \\ \omega(1) & \text{if } p = \omega\left(n^{-2/(k-1)}\right) \end{cases}$$

- **Chebyshev**: $\Pr(X = 0) \leq \Pr(|X - \mathbb{E}[X]| \geq \mathbb{E}[X]) \leq \dfrac{\mathbf{Var}[X]}{\mathbb{E}[X]^2} \leq \dfrac{1}{\mathbb{E}[X]} + \dfrac{\sum_{S \neq T} \mathbb{E}[I_S I_T]}{\mathbb{E}[X]^2}$

$$= O\left(n^{-k} p^{-\binom{k}{2}}\right) + O\left(\sum_{\ell=2}^{k-1} n^{-\ell} p^{-\binom{\ell}{2}}\right) = O\left(\sum_{\ell=2}^{k} n^{-\ell} p^{-\binom{\ell}{2}}\right)$$

$$= o(1) \text{ if } p = \omega\left(n^{2/(1-k)}\right)$$

- $\implies \Pr(X \geq 1) \geq 1 - o(1)$

A "Threshold Behavior" in Random Graphs
(Erdős–Rényi 1960)

- Fix a constant integer $k \geq 3$.
- Let $G \sim G(n, p)$, as $n \to \infty$:

$$\Pr(G \text{ contains a } K_k) = \begin{cases} o(1) & \text{if } p = o\left(n^{-2/(k-1)}\right) \\ 1 - o(1) & \text{if } p = \omega\left(n^{-2/(k-1)}\right) \end{cases}$$

- For $H(V, E)$ with $k = |V|$, $m = |E|$ s.t. every subgraph of H has density $\leq m/k$:

$$\Pr(G \text{ contains a subgraph } H) = \begin{cases} o(1) & \text{if } p = o\left(n^{-k/m}\right) \\ 1 - o(1) & \text{if } p = \omega\left(n^{-k/m}\right) \end{cases}$$

Weierstrass Approximation Theorem

(魏尔施特拉斯逼近定理)

- <u>Weierstrass Approximation Theorem</u>: Let $f : [0,1] \to [0,1]$ be a continuous function. For any $\epsilon > 0$, there exists a polynomial p such that

$$\sup_{x \in [0,1]} |p(x) - f(x)| \leq \epsilon$$

- **Proof**: Let integer n be sufficiently large (to be fixed later).

  For $x \in [0,1]$, let $X \sim \frac{1}{n}\text{Bin}(n, x)$. Define polynomial p on $x \in [0,1]$ to be:

$$p(x) = \mathbb{E}\left[f(X)\right] = \sum_{k=0}^{n} f\left(\frac{k}{n}\right)\binom{n}{k} x^k (1-x)^{n-k}$$

Let $f : [0,1] \to [0,1]$ be continuous. For $x \in [0,1]$, let $X \sim \frac{1}{n}\text{Bin}(n, x)$, and:

$$p(x) = \mathbb{E}\left[f(X)\right] = \sum_{k=0}^{n} f\left(\frac{k}{n}\right)\binom{n}{k} x^k (1-x)^{n-k}$$

$$|p(x) - f(x)| = \left|\mathbb{E}\left[f(X) - f(x)\right]\right| \leq \mathbb{E}\left[\left|f(X) - f(x)\right|\right]$$

(f is continuous on $[0,1] \implies \exists \delta > 0$ s.t. $|f(x) - f(y)| \leq \epsilon/2$ for all $|x - y| \leq \delta$)

$$= \mathbb{E}\left[\left|f(X) - f(x)\right| \mid \left|X - x\right| \leq \delta\right] \cdot \Pr\left(\left|X - x\right| \leq \delta\right)$$

$$+ \mathbb{E}\left[\left|f(X) - f(x)\right| \mid \left|X - x\right| > \delta\right] \cdot \Pr\left(\left|X - x\right| > \delta\right)$$

$$\leq \mathbb{E}\left[\epsilon/2\right] + \left|1 - 0\right| \cdot \Pr\left(\left|X - x\right| > \delta\right) \quad \leq \frac{\epsilon}{2} + \frac{x(1-x)}{n\delta^2}$$ (Chebyshev)

$$\leq \frac{\epsilon}{2} + \frac{1}{4n\delta^2} \quad \leq \epsilon$$ if we choose $n \geq \frac{1}{2\epsilon\delta^2}$

Weierstrass Approximation Theorem

(魏尔施特拉斯逼近定理)

- <u>Weierstrass Approximation Theorem</u>: Let $f : [0,1] \to [0,1]$ be a continuous function. For any $\epsilon > 0$, there exists a polynomial p such that

$$\sup_{x \in [0,1]} |p(x) - f(x)| \leq \epsilon$$

- **Proof**: By continuity, $\exists \delta > 0$ s.t. $|f(x) - f(y)| \leq \epsilon/2$ if $|x - y| \leq \delta$.

  Let $n \geq 1/(2\epsilon\delta^2)$ be any integer. For $x \in [0,1]$, let $X \sim \frac{1}{n}\text{Bin}(n, x)$, and:

$$p(x) = \mathbb{E}\left[f(X)\right] = \sum_{k=0}^{n} f\left(\frac{k}{n}\right)\binom{n}{k} x^k (1-x)^{n-k}$$

  For any $x \in [0,1]$, it holds that $|p(x) - f(x)| \leq \epsilon$.

Higher Moments





Skewness (偏度)

- The <u>skewness</u> (偏度) of a random variable X with expectation $\mu = \mathbb{E}[X]$ and standard deviation $\sigma = \sqrt{\mathbf{Var}[X]}$ is defined by

$$\mathrm{Skew}[X] = \mathbb{E}\left[\left(\frac{X-\mu}{\sigma}\right)^3\right] = \frac{\mathbb{E}[(X-\mu)^3]}{\sigma^3}$$

standardized moment (of degree 3)



Kurtosis (峰度)

- The <u>**kurtosis**</u> (峰度) of a random variable X with expectation $\mu = \mathbb{E}[X]$ and standard deviation $\sigma = \sqrt{\mathbf{Var}[X]}$ is defined by

$$\mathrm{Kurt}[X] = \mathbb{E}\left[\left(\frac{X-\mu}{\sigma}\right)^4\right] = \frac{\mathbb{E}[(X-\mu)^4]}{\sigma^4}$$

standardized moment (of degree 4)



The kth Moment Method

- Let X be a random variable with $\mathbb{E}[X] = \mu$. For any $C > 1$ and integer $k \geq 1$

$$\Pr\left(|X - \mu| \geq C \cdot \mathbb{E}\left[|X - \mu|^k \right]^{\frac{1}{k}} \right) \leq \frac{1}{C^k}$$

- **Proof**: Apply Markov's inequality to $Z = |X - \mu|^k$.

The Moment Problem

- Do moments $m_k = \mathbb{E}[X^k]$, $\forall k \geq 1$, uniquely identify the distribution of X?
- If X takes values from a **finite** set $\{x_1, \ldots, x_n\}$,

  then solving the Vandermonde system:

$$\begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ x_1^2 & x_2^2 & \cdots & x_n^2 \\ \vdots & \vdots & \ddots & \vdots \\ x_1^n & x_2^n & \cdots & x_n^n \end{bmatrix} \begin{bmatrix} p_1 \\ p_2 \\ \vdots \\ p_n \end{bmatrix} = \begin{bmatrix} m_1 \\ m_2 \\ \vdots \\ m_n \end{bmatrix}$$

  can recover the *pmf* $p_i = p_X(x_i)$

The Moment Problem

- Do moments $m_k = \mathbb{E}[X^k]$, $\forall k \geq 1$, uniquely identify the distribution of X?
  - If $\mathbb{E}[X^k] = \mathbb{E}[Y^k]$ for all $k \geq 1$, are X and Y always identically distributed?
- If X and Y have the same <u>**moment generating function (MGF)**</u>

$$M_X(t) = \mathbb{E}[\mathrm{e}^{tX}] \ = \sum_{k \geq 0} \frac{t^k \mathbb{E}[X^k]}{k!}$$

  then X and Y are identically distributed.
- The MGF $M_X(t)$ is convergent if the sequence $\mathbb{E}[X^k]$ does not grow too fast.

Moment Generating Function

- The <u>**moment generating function**</u> (<u>**MGF**</u>) of a random variable X is

$$M_X(t) = \mathbb{E}[\mathrm{e}^{tX}]$$

- Expand e^{tX} by Maclaurin series: $M_X(t) = \mathbb{E}[\mathrm{e}^{tX}] \ = \sum_{k \geq 0} \frac{t^k \mathbb{E}[X^k]}{k!}$

- In fact, the kth moment is given by $\mathbb{E}[X^k] = M_X^{(k)}(0)$ where $M_X^{(k)}(0)$ is the kth derivative of $M_X(t)$ evaluated at $t = 0$

- If $M_X(t) = M_Y(t)$ for all $t \in [-\delta, \delta]$ for some $\delta > 0$, then X, Y are identically distributed.

Poisson Tails



- A Poisson random variable $X \sim \text{Pois}(\lambda)$:

$$p_X(k) = \Pr(X = k) = \mathrm{e}^{-\lambda}\frac{\lambda^k}{k!}, \quad k = 0,1,2,\ldots$$

- Tail inequalities for $X \sim \text{Pois}(\lambda)$:

$$k > \lambda \implies \Pr(X \geq k) \leq \mathrm{e}^{-\lambda}\left(\frac{\mathrm{e}\lambda}{k}\right)^k$$

$$k < \lambda \implies \Pr(X \leq k) \leq \mathrm{e}^{-\lambda}\left(\frac{\mathrm{e}\lambda}{k}\right)^k$$

- **Chebyshev:** $\Pr\left[|X - \lambda| \geq |k - \lambda|\right] \leq \dfrac{\lambda}{(k-\lambda)^2}$

MGF of Poisson Random Variable

- A Poisson random variable $X \sim \text{Pois}(\lambda)$:

$$p_X(k) = \Pr(X = k) = \mathrm{e}^{-\lambda}\frac{\lambda^k}{k!}, \quad k = 0,1,2,\ldots$$

- **MGF**: $M_X(t) = \mathbb{E}\left[\mathrm{e}^{tX}\right] = \sum_{k=0}^{\infty} \frac{\mathrm{e}^{-\lambda}\lambda^k}{k!}\mathrm{e}^{tk}$

$$= \mathrm{e}^{\lambda(\mathrm{e}^t - 1)} \sum_{k=0}^{\infty} \frac{\mathrm{e}^{-\lambda \mathrm{e}^t}(\lambda \mathrm{e}^t)^k}{k!}$$

$$= \mathrm{e}^{\lambda(\mathrm{e}^t - 1)}$$

Cramér-Chernoff Method (Chernoff Bound)

- MGF for $X \sim$ Pois(λ):

$$M_X(t) = \mathbb{E}\left[\mathrm{e}^{tX}\right] = \mathrm{e}^{\lambda(\mathrm{e}^t - 1)}$$

- Upper tail: $\Pr(X \geq k) = \Pr\left(\mathrm{e}^{tX} \geq \mathrm{e}^{tk}\right)$ (for all $t > 0$)

(Markov's Ineq.) $$\leq \frac{\mathbb{E}\left[\mathrm{e}^{tX}\right]}{\mathrm{e}^{tk}} = \mathrm{e}^{\lambda(\mathrm{e}^t - 1) - tk}$$

(set $t = \ln(k/\lambda) > 0$) $$= \mathrm{e}^{-\lambda}\left(\frac{\mathrm{e}\lambda}{k}\right)^k$$

Cramér-Chernoff Method (Chernoff Bound)

- MGF for $X \sim$ Pois(λ):

$$M_X(t) = \mathbb{E}\left[\mathrm{e}^{tX}\right] = \mathrm{e}^{\lambda(\mathrm{e}^t - 1)}$$

- Lower tail: $\Pr(X \leq k) = \Pr\left(\mathrm{e}^{tX} \geq \mathrm{e}^{tk}\right)$ (for all $t < 0$)

(Markov's Ineq.) $\leq \dfrac{\mathbb{E}\left[\mathrm{e}^{tX}\right]}{\mathrm{e}^{tk}} = \mathrm{e}^{\lambda(\mathrm{e}^t - 1) - tk}$

(set $t = \ln(k/\lambda) < 0$) $= \mathrm{e}^{-\lambda}\left(\dfrac{\mathrm{e}\lambda}{k}\right)^k$